

Going to the bathroom too often?

We can help.

Loss of bladder and bowel control are extremely common.¹⁻⁴ Choose a more personalized approach and get more control.

If you are experiencing any of the following:

- Uncontrollable urges to go to the bathroom
- Frequent accidents resulting in wearing pads
- Getting up multiple times at night
- Not emptying your bladder all the way

Join us for a free, anonymous, virtual educational event

- Learn about symptoms, conditions, and causes
- Understand your treatment options
- Understand next steps to gain more freedom, more control, and more life

Date:

Time:

Registration:

Speaker(s):



Co-sponsored by

Medtronic



1. Stewart WF, et al. Prevalence and burden of overactive bladder in the United States. *World J Urol.* 2003 May;20(6):327-336.
2. US Census Bureau 2020. US adult and under-age-18 populations: 2020 census. <https://www.census.gov/library/visualizations/interactive/adult-and-under-the-age-of-18-populations-2020-census.html>. Accessed June 20, 2022.
3. Ditah I, Devaki P, Luma HN, et al. Prevalence, trends, and risk factors for fecal incontinence in United States adults, 2005-2010. *Clin Gastroenterol Hepatol.* 2014;12:636-643.
4. Whitehead WE, Borrud L, Goode PS, et al. Fecal Incontinence in US adults: epidemiology and risk factors. *Gastroenterology.* 2009;137(2):512-517.

